

The facial weak order in hyperplane arrangements

Aram Dermenjian^{1,3}

Christophe Hohlweg¹, Thomas McConville² and Vincent Pilaud³

¹Université du Québec à Montréal (UQAM)

²Mathematical Sciences Research Institute (MSRI)

³École Polytechnique (LIX)

10 May 2019

On this day in 1847 Wilhelm Killing was born.


Outline

- Arranging hyperplanes.
- The facial weak order and its 1, 2, 3, 4 (!) definitions.
- Yeah, but is it a lattice?
- Some other properties.

History and Background - Hyperplanes

- $(V, \langle \cdot, \cdot \rangle)$ - n -dim real Euclidean vector space.
- A *hyperplane* H_i is codim 1 subspace of V with normal e_i .




Example



History and Background - Arrangements


- A *hyperplane arrangement* is $\mathcal{A} = \{H_1, H_2, \dots, H_k\}$.
- \mathcal{A} is *central* if $\{0\} \subseteq \bigcap \mathcal{A}$.
- Central \mathcal{A} is *essential* if $\{0\} = \bigcap \mathcal{A}$.

Example



History and Background - Arrangements

- *Regions* \mathcal{R}_A - connected components of V without \mathcal{A} .
- *Faces* \mathcal{F}_A - intersections of closures of some regions.




History and Background - (Partial) Orders

- *Lattice* - poset where every two elements have a *meet* (greatest lower bound) and *join* (least upper bound).

Example


The lattice $(\mathbb{N}, |)$ where $a \leq b \Leftrightarrow a | b$.

- meet - greatest common divisor
- join - least common multiple




History and Background - Poset of regions

- *Base region* $B \in \mathcal{R}_A$ - some fixed region
- *Separation set for* $R \in \mathcal{R}_A$
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$




History and Background - Poset of regions

- *Base region* $B \in \mathcal{R}_A$ - some fixed region
- *Separation set for* $R \in \mathcal{R}_A$
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$




History and Background - Poset of regions

- *Base region* $B \in \mathcal{R}_A$ - some fixed region
- *Separation set for* $R \in \mathcal{R}_A$
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$




History and Background - Poset of regions

- *Base region* $B \in \mathcal{R}_A$ - some fixed region
- *Separation set for* $R \in \mathcal{R}_A$
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$



History and Background - Poset of regions



- *Base region* $B \in \mathcal{R}_A$ - some fixed region
- *Separation set for* $R \in \mathcal{R}_A$
 $S(R) := \{H \in \mathcal{A} \mid H \text{ separates } R \text{ from } B\}$
- *Poset of Regions* $\text{PR}(\mathcal{A}, B)$ where
 $R \leq_{\text{PR}} R' \Leftrightarrow S(R) \subseteq S(R')$



History and Background - Poset of regions

- A region R is *simplicial* if normal vectors for boundary hyperplanes are linearly independent.
- \mathcal{A} is *simplicial* if all \mathcal{R}_A simplicial.

Example






History and Background - Poset of regions

Theorem (Björner, Edelman, Ziegler '90)

If \mathcal{A} is simplicial then $\text{PR}(\mathcal{A}, B)$ is a lattice for any $B \in \mathcal{R}_{\mathcal{A}}$. If $\text{PR}(\mathcal{A}, B)$ is a lattice then B is simplicial.

Example






History and Background - Poset of regions

Theorem (Björner, Edelman, Ziegler '90)

If \mathcal{A} is simplicial then $\text{PR}(\mathcal{A}, B)$ is a lattice for any $B \in \mathcal{R}_{\mathcal{A}}$. If $\text{PR}(\mathcal{A}, B)$ is a lattice then B is simplicial.

Example



Coxeter Arrangements

Example

A *Coxeter arrangement* is the hyperplane arrangement associated to a Coxeter group.

Coxeter Groups		Hyperplane Arrangements
Reflecting hyperplanes	\leftrightarrow	Hyperplane arrangement
Root system	\leftrightarrow	Normals to hyperplanes
Inversion sets	\leftrightarrow	Separation sets
Weak order	\leftrightarrow	Poset of regions

Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order of Coxeter groups to an order on all the faces of its associated arrangement for type A (aka Braid arrangement).
- In 2006, Palacios and Ronco extended this new order to Coxeter groups of all types using cover relations.
- In 2016, D, Hohlweg and Pilaud showed this extension has a global equivalent and produces a lattice in Coxeter arrangements.


Motivation

- In 2001, Krob, Latapy, Novelli, Phan, and Schwer extended the weak order of Coxeter groups to an order on all the faces of its associated arrangement for type A (aka Braid arrangement).
- In 2006, Palacios and Ronco extended this new order to Coxeter groups of all types using cover relations.
- In 2016, D, Hohlweg and Pilaud showed this extension has a global equivalent and produces a lattice in Coxeter arrangements.
- Questions: Can we extend this to hyperplane arrangements? Can we find both local and global definitions? When do we actually get a lattice?

Facial Intervals

Proposition (Björner, Las Vergas, Sturmfels, White, Ziegler '93)

Let \mathcal{A} be central with base region B . For every $F \in \mathcal{F}_{\mathcal{A}}$ there is a unique interval $[m_F, M_F]$ in $\text{PR}(\mathcal{A}, B)$ such that

$$[m_F, M_F] = \{R \in \mathcal{R}_{\mathcal{A}} \mid F \subseteq \overline{R}\}$$


Facial Weak Order



Let \mathcal{A} be a central hyperplane arrangement and B a base region in $\mathcal{R}_{\mathcal{A}}$.

Definition



The *facial weak order* is the order $\text{FW}(\mathcal{A}, B)$ on $\mathcal{F}_{\mathcal{A}}$ where for $F, G \in \mathcal{F}_{\mathcal{A}}$:

$$F \leq G \Leftrightarrow m_F \leq_{\text{PR}} m_G \text{ and } M_F \leq_{\text{PR}} M_G$$



Facial Weak Order - Example





Facial Weak Order - Example





Facial Weak Order - Example





Facial Weak Order - Example





Facial Weak Order - Example



Facial Weak Order - Example



Facial Weak Order - Example





Cover Relations

Proposition (D., Hohlweg, McConville, Pilaud, '19+)

For $F, G \in \mathcal{F}_A$ if

1. $F \leq G$ in $\text{FW}(\mathcal{A}, B)$
2. $|\dim(F) - \dim(G)| = 1$
3. $F \subseteq G$ or $G \subseteq F$

then $F \lessdot G$.




Covectors

- *covector* - a vector in $\{-, 0, +\}^A$ with signs relative to hyperplanes.
- $\mathcal{L} \subseteq \{-, 0, +\}^A$ - set of covectors

Example

$$F_4 \leftrightarrow (+, 0, -) \quad F_4(H_1) = +; F_4(H_2) = 0; F_4(H_3) = -$$




Covectors

- *covector* - a vector in $\{-, 0, +\}^A$ with signs relative to hyperplanes.
- $\mathcal{L} \subseteq \{-, 0, +\}^A$ - set of covectors

Example

$$F_4 \leftrightarrow (+, 0, -) \quad F_4(H_1) = +; F_4(H_2) = 0; F_4(H_3) = -$$



Covector operations

For $X, Y \in \mathcal{L} \subseteq \{-, 0, +\}^{\mathcal{A}}$

■ *Composition*: $(X \circ Y)(H) = \begin{cases} Y(H) & \text{if } X(H) = 0 \\ X(H) & \text{otherwise} \end{cases}$

■ *Reorientation*: $(X_{-Y})(H) = \begin{cases} -X(H) & \text{if } Y(H) = 0 \\ X(H) & \text{otherwise} \end{cases}$

★ For $F \in \mathcal{F}_A$, $[m_F, M_F] = [F \circ B, F \circ -B]$

Example

Let $\mathcal{A} = \{H_1, H_2, H_3, H_4, H_5\}$.

$$X = (-, 0, +, +, 0) \quad Y = (0, 0, -, 0, +)$$

Then

$$X \circ Y = (-, 0, +, +, +) \quad X_{-Y} = (+, 0, +, -, 0)$$





Covector Definition

Definition

For $X, Y \in \mathcal{L}$:

$$X \leq_{\mathcal{L}} Y \Leftrightarrow X(H) \geq Y(H) \quad \forall H \text{ with } - < 0 < +$$



Zonotopes

- *Zonotope* $Z_{\mathcal{A}}$ is the convex polytope:

$$Z_{\mathcal{A}} := \left\{ v \in V \mid v = \sum_{i=1}^k \lambda_i e_i, \text{ such that } |\lambda_i| \leq 1 \text{ for all } i \right\}$$


Theorem (Edelman '84, McMullen '71)

There is a bijection between $\mathcal{F}_{\mathcal{A}}$ and the nonempty faces of $Z_{\mathcal{A}}$ given by the map


$$\tau(F) = \left\{ v \in V \mid v = \sum_{F(H_i)=0} \lambda_i e_i + \sum_{F(H_j) \neq 0} \mu_j e_j \right\}$$

where $|\lambda_i| \leq 1$ for all i and $\mu_j = F(H_j)$


Zonotope - Construction example




Zonotope - Construction example




Zonotope - Construction example




Zonotope - Construction example



Zonotope - Construction example



Zonotope - Construction example





Root inversion sets

■ roots $\Phi_{\mathcal{A}} := \{\pm e_1, \pm e_2, \dots, \pm e_k\}$

■ root inversion set

$$\mathbf{R}(F) := \{e \in \Phi_{\mathcal{A}} \mid \langle x, e \rangle \leq 0 \text{ for some } x \in F\}.$$



Equivalent definitions

Theorem (D., Hohlweg, McConville, Pilaud '19+)





For $F, G \in \mathcal{F}_{\mathcal{A}}$ the following are equivalent:

- $m_F \leq_{\text{PR}} m_G$ and $M_F \leq_{\text{PR}} M_G$ in poset of regions $\text{PR}(\mathcal{A}, B)$.
- There exists a chain of covers in $\text{FW}(\mathcal{A}, B)$ such that





$$F = F_1 \triangleleft F_2 \triangleleft \cdots \triangleleft F_n = G$$

- $F \leq_{\mathcal{L}} G$ in terms of covectors ($F(H) \geq G(H) \forall H \in \mathcal{A}$)
- $\mathbf{R}(F) \setminus \mathbf{R}(G) \subseteq \Phi_{\mathcal{A}}^-$ and $\mathbf{R}(G) \setminus \mathbf{R}(F) \subseteq \Phi_{\mathcal{A}}^+$.


Equivalence for type A_2 Coxeter arrangement




Equivalence for type A_2 Coxeter arrangement







Equivalence for type A_2 Coxeter arrangement








Equivalence for type A_2 Coxeter arrangement




Equivalence for type A_2 Coxeter arrangement







Equivalence for type A_2 Coxeter arrangement



Equivalence for type A_2 Coxeter arrangement



Equivalence for type A_2 Coxeter arrangement



Facial weak order lattice

Theorem (D., Hohlweg, McConville, Pilaud '19+)

The facial weak order $\text{FW}(\mathcal{A}, B)$ is a lattice when \mathcal{A} is simplicial.

Corollary (D., Hohlweg, McConville, Pilaud '19+)

The lattice of regions is a sublattice of the facial weak order lattice when \mathcal{A} is simplicial.

Lattice proof - Joins

Proof uses two key components :


Lemma (Björner, Edelman, Ziegler '90)

1: If L is a finite, bounded poset such that $x \vee y$ exists whenever x and y both cover some $z \in L$, then L is a lattice.


2: Cover relation: $Z \triangleleft X$ iff $Z \leq X$, $|\dim X - \dim Z| = 1$ and $X \subseteq Z$ or $Z \subseteq X$. Then $Z \triangleleft X$ and $Z \triangleleft Y$ gives three cases:

1. $X \cup Y \subseteq Z$ and $\dim X = \dim Y = \dim Z - 1$,
2. $Z \subseteq X \cap Y$ and $\dim X = \dim Y = \dim Z + 1$, and
3. $X \subseteq Z \subseteq Y$ and $\dim X = \dim Z - 1 = \dim Y - 2$.


$X \cup Y \subseteq Z$ and $\dim X = \dim Y = \dim Z - 1$




$X \cup Y \subseteq Z$ and $\dim X = \dim Y = \dim Z - 1$




$Z \subseteq X \cap Y$ and $\dim X = \dim Y = \dim Z + 1$




$Z \subseteq X \cap Y$ and $\dim X = \dim Y = \dim Z + 1$




$X \subseteq Z \subseteq Y$ and $\dim X = \dim Z - 1 = \dim Y - 2$




$X \subseteq Z \subseteq Y$ and $\dim X = \dim Z - 1 = \dim Y - 2$




$X \subseteq Z \subseteq Y$ and $\dim X = \dim Z - 1 = \dim Y - 2$



$X \subseteq Z \subseteq Y$ and $\dim X = \dim Z - 1 = \dim Y - 2$



Example: B_3 Coxeter arrangement



Properties of the facial weak order

- The *dual* of a poset P is the poset P^{op} where $x \leq y$ in P iff $y \leq x$ in P^{op} . A poset is *self-dual* if $P \cong P^{op}$.
- A lattice is *semi-distributive* if $x \vee y = x \vee z$ implies $x \vee y = x \vee (y \wedge z)$ and similarly for the meets.

Theorem (D., Hohlweg, McConville, Pilaud '19+)

The facial weak order $\text{FW}(\mathcal{A}, B)$ is self-dual. If furthermore, \mathcal{A} is simplicial, $\text{FW}(\mathcal{A}, B)$ is a semi-distributive lattice.

Join-irreducible elements

- An element is *join-irreducible* if and only if it covers exactly one element.

Proposition (D., Hohlweg, McConville, Pilaud '19+)

If \mathcal{A} is simplicial and F a face with facial interval $[m_F, M_F]$. Then F is join-irreducible in $\text{FW}(\mathcal{A}, B)$ if and only if M_F is join-irreducible in $\text{PR}(\mathcal{A}, B)$ and $\text{codim}(F) \in \{0, 1\}$

Möbius function

Recall that the Möbius function is given by:

$$\mu(x, y) = \begin{cases} 1 & \text{if } x = y \\ -\sum_{x \leq z < y} \mu(x, z) & \text{if } x < y \\ 0 & \text{otherwise} \end{cases}$$


Proposition (D., Hohlweg, McConville, Pilaud '19+)

Let X and Y be faces such that $X \leq Y$ and let $Z = X \cap Y$.

$$\mu(X, Y) = \begin{cases} (-1)^{\text{rk}(X) + \text{rk}(Y)} & \text{if } X \leq Z \leq Y \text{ and } Z = X_{-Z} \cap Y \\ 0 & \text{otherwise} \end{cases}$$

Further Works

- Can we explicitly state the join/meet of two elements?
- When is the facial weak order congruence uniform?
- Can we generalize this to polytopes?



Thank you!